



STUDIES ON SEED PATHOLOGY OF CERTAIN VEGETABLES

DISSERTATION SUBMITTED FOR THE DEGREE OF
Master of Philosophy
IN
BOTANY

BY
MOHD. ABUL FAZAL

DEPARTMENT OF BOTANY
ALIGARH MUSLIM UNIVERSITY
ALIGARH (INDIA)

May, 1987



DS1130

DEDICATED TO
LATE DR. DHIRENDRA PRAKASH

C O N T E N T S

PAGE NO.

1.	INTRODUCTION	.1..
2.	REVIEW OF LITERATURE	.6..
3.	MATERIALS AND METHODS	.32.
4.	PLAN OF WORK	.36.
5.	REFERENCES	.37.

=====

I N T R O D U C T I O N

=====

INTRODUCTION

Vegetables are natural sources of proteins, carbohydrates, vitamins and minerals, therefore, occupy an important position in the human diet, particularly in India where major population is vegetarian. Diseases and pests are responsible for heavy damages to vegetable every year (Walker, 1952 and Prem Nath, 1976). Amongst the various diseases, seed-borne diseases are of no less importance. Considerable work has been done on the seed pathology of vegetable seeds (Neergard, 1977). ✓ But in all these studies one thing emerges that amongst various types of pathogens reported from seed, fungi constitute a major portion causing seed rots, seedling rot i.e. pre-and post-emergence damping off losses at various stages of plant growth.

Infection of vegetable to the extent of 50 percent has been reported in many areas of the world. Cramer (1967) pointed out that although statistical data on world wide losses are not available for the majority of vegetables, but it is felt that losses are enormous. In the U.S.A. annual average loss of bean (Phaseolus vulgaris) was 20 percent of which 16 to 17 percent was due to seed borne infection (Anon, 1965) and of these one third was attributed to Fusarium solani f. sp. phaseoli alone. ✓ In 1939 Fusarium

root rot caused 95 percent losses of beans in New South Wales, Australia (Anon,1941). In South Western Colarado 84 percent yield reduction was recorded to vegetable crops and not a single field sample was found free of the disease (Keenan et al.,1974). In temperate countries Collectotrichum lindemuthianum the causal agent of bean anthracnose, a seed-borne fungus, has been reported to cause tremendous loss to beans. Masurat et al.(1966) reported that the damage of beans to the extent of five percent was due to anthracnose during 1965 in East Germany. While Chang (1960) observed 50 - 54 in U.S.S.R. Losses upto 50 percent have been reported in the U.S.A. (Zaumeyer and Thomas,1957) and in recent years losses have increased alarmingly. In Western Nebraska the average loss in dry beans was 6 to 19 percent in different years from 1970 - 72 (Kerr and Steadman,1974) while in New York it was 10 to 20 percent (Lumsden et al., 1974). Leaf spot causing pathogen Ascochyta spp. of beans and kidney beans was also found seed-borne and was reported causing considerable losses (Ondrej,1971). Watanabe (1972) isolated virulent Macrophomina phaseoli from commercial kidney bean seed lots. Where as, seed rot and root rot fungi Fusarium solani Phylosticta phaseolina were recorded from seeds of kidney bean (Gupta and Saharan,1973). Ascochyta infected bean seeds when sowed produced lesions on plants to the extent of 2 to 15 percents. (Hewett,1973). Ellis et al (1976) reported a correlation between seed mycoflora

and percentage germination of bean seed. Harrison (1978) discussed the role of seed-borne inoculum in the epidemiology of Botrytis fabae on field beans.

Ascochyta complex, Fusarium wilt and bacterial blight caused by Ascochyta spp., Fusarium oxysporum f. sp. pisi and Pseudomonas pisi on pea are responsible for heavy loss every year. In the U.S.A. average annual losses of the national production of green peas for the period 1951-1960 (Anon, 1965) were estimated to 2 percent for Ascochyta complex, 2 percent for Fusarium wilt and 1 percent for bacterial blight. In Canada, Wallen (1965) found experimentally that A. pisi under certain conditions caused yield reduction to 11 percent, A. pinodella 25 percent and A. pinodes 45 percent. Teranishi and Namekata (1972) observed wilting in pea plant by Fusarium oxysporum f. sp. pisi.

Cicer arietinum is badly attacked by Ascochyta rabiei a causal agent of Ascochyta blight or leaf blight and some times Sclerotinia sclerotiorum, stem and crown blight which are seed-borne in nature. In 1969 A. rabiei was found to cause severe dieback of foliage, and yields were so much reduced that it became a limiting factor in cultivation (Kaiser, 1970). Halfon-Meiri (1970) observed a reduction in seed weight by 40 percent due to Ascochyta. Similarly losses ranging from 15 - 83 percent due to this disease has also been observed (Askerov, 1968; Morall and McKenzie, 1974).

Amongst foliage and green vegetables the importance of cabbage and cauliflower can not be under estimated. The seeds of these vegetables are not free from fungi which are seed-borne such as Phoma lingam, dry rot, black leg and Xanthmmones compestris black rot and Alternaria brassicicola leaf spot. These are responsible not only for lowering the quality of seed but also decreasing the yield of seed. Giessmann and Daebler (1973) reported that Phoma lingam was a seed-borne pathogen on cabbage and was responsible for heavy losses. Heavy toll of various foliage vegetable crops due to seed-borne Alternaria has been reported from many parts of the world. (Holtzhausen, 1978 and Wu et al., 1979). Singh et al. (1970) reported that Helminthosporium spiciferum and Curvularia lunata and leaf spot pathogens of egg plant are seed-borne and are responsible for heavy losses every year in India. Losses to the extent of 25% on okra due to Rhizoctonia bataticola and Choenophora cucurbitarum have been observed by Goel and Mehrotra (1973) Tai Luang Haun and Musa Bin Jamail (1975) and Geol and Mehrotra (1978). Fusarium oxysporum f. sp. spinaceae and Phoma vallerianellae are seed-borne (Bassi et al., 1978) and take heavy toll of the crop every year in France (Vegh et al., 1978).

It is evident from the above that seed-borne fungi are responsible of causing severe losses to vegetables in field by reducing the germination ability, resulting in damping off, seedling rot etc. Ultimately the crop is damaged and the yield is reduced. In India although considerable work has been done on seed mycoflora of vegetable seeds but the information with regard to vegetable seeds is still far from satisfactory. Therefore, in the present investigation an attempt will be made to determine seed mycoflora of selected vegetable seeds such as peas, cabbage, cauliflower etc. and to work out control measures to minimise the losses.

=====

R E V I E W O F L I T E R A T U R E

=====

REVIEW OF LITERATURE

Studies on seed pathology infact can be dated back to 1637 when Remnant reported that seeds carry disease causing agents. The pioneer work of Hellwig (1699) with Claviceps purpurea (Fr.) Tul. on rye, Micheli (1723) with Orbanche minor L. on beans and Tillet (1755) with Tilletia sp. on wheat established that disease causing agents are carried by seeds. Frank (1883) demonstrated that the bean anthracnose fungus Colletotrichum lindemuthianum was seed-borne in nature.

The first bacterial pathogen to be recorded as seed-borne was Xanthomonas phaseoli causing common bean blight (Orton, 1931).

Needham (1745) for the first time reported seed-borne nature of nematode Anguina tritici on wheat. This was followed by Kuhn (1857) who indicated seed-borne nature of Ditylenchus dipsaci.

Amongst the viral diseases lima bean mosaic was for first time suspected by McClintock (1916) as seed-borne. Subsequently several viral diseases have been reported seed-borne (Leach, 1940; Carter, 1966; Narayanaswamy and Jagannathan, 1975).

Realising the importance of losses due to seed-borne diseases the first official seed testing station was established in Saxony in 1869 to determine the purity and germination capacity of seeds (Malone and Muskett, 1964). However, before the establishment of seed testing station, the Switzerland government passed a law requiring the examination of seeds (Porter, 1949). This was followed by realisation of losses due to seed-borne diseases in other countries. As a result of this more and more seed testing stations were established.

Muskett (1950) pointing out the significance of seed-borne diseases observed several fungi associated with seeds. Extensive studies on external seed-borne fungal flora of wheat were made by Christensen (1951) and of rice by Cherewick (1954). Kommendahl et al. (1955) while investigating the seed mycoflora observed Alternaria spp. and Colletotrichum spp. to be very frequent on seeds of flax. Schmidt (1955) reported 50 different species of fungi from certain agriculturally important crops. Detailed studies on seed-borne fungi on wheat (Russel 1956; De-Tempe, 1964) pine seed (Gibson 1957) and small grains (Crosier, 1961); guar, beans (Jain and Patil, 1969), ground nut (Joffe 1969; Gupta and Chohan, 1971), Cucumbers, bottlegourd (Khandelwal and Prasad, 1970), Cauliflower, bean and pea (Sekhon and Shivapuri, 1971; Haware, 1971) have been made. De-Tempe (1964a) pointed out that Fusarium sp. was seed-borne on cereals

of temperate countries. Phoma betae, Alternaria tenuis and Fusarium sp. were more frequent on sugarbeet seed (Gambogi, 1964) and Alternaria dauci and Stemphylium redicinum on carrot (Hewett, 1964); Ascochyta fabae on pea (Hewett, 1966) A. pisi, A. pinodella and Macrophomina phaseoli on pea (Bedlam, 1985) A. rabei on chickpea (Halfon-Meiri 1970; Maden et al., 1975) was also found to be seed-borne. Seed-borne nature of the pathogen was also found to be for haloblight and anthracnose of bean. Watanabe (1972) established the seed-borne nature of large number of fungi on kidney bean, including some very serious. Aspergillus flavus was very frequently observed on the soybean et al. (Dhingra/1973); Rhizoctonia bataticola on okra (Goel and Mehrotra, 1973). Scott and Wenham (1973) intercepted Alternaria radicina and A. dauci from the seeds of carrot imported to Newzealand. Tylkowska (1984) observed prevalence of large number of fungi including A. radicina and A. dauci on seven cultivars of carrot, while A. alternata was reported to be a serious pathogen of snap bean by Gomes and Dhingra (1983). In India Phoma betae, Cercospora bataticola and Verticillium spp. were intercepted in exotic lots of sugar beet seeds (Lambat et al., 1974) while Singh et al. (1974) isolated 17 genera of fungi. A comprehensive list of fungi associated with pea seed has been made by Bedi and Kapur (1974), while that of pepper and egg plant by Kar'kova et al. (1974); of cruciferous vegetable crops by Holtzhausen and

Knox-Davles (1974) and Toskova et al.(1975); of carrot by Mirkova (1976); of rape seed by Mills and Bollen (1976); of bottle-gourd by Dwivedi and Tandon (1976); of pulse crops by Suhag and Suryanarayan (1976); of bean by Lasca (1978), Kanapathipillai (1982) and Tylkowska (1984); of cruciferous crops by Tahvonen (1979) and Kanapathipillai and Hashim (1981); of soybean by Zad (1979) and Kumar et al.(1980); of cowpea by Kumari and Karan (1981); Zaidi and Prakash (1983) and Baross et al. (1985); of gram by D'Ercole and Sportelli (1982); of *Raphanus sativus* by Thakur et al.(1982) and of cucurbits by Komoto and Kimura (1982). Abdalla (1974) reported that although Aspergillus flavus and A.niger were frequently found in all the samples of seeds of ground nut but its frequency was very high in those having fractured shells. Pflieger and Herman (1975) observed Aspergillus flavus externally from pea seeds but it did not infect the embryo. Lorenzini et al.(1975) found Fusarium oxysporum f.sp. pisi isolated from seeds pathogenic and responsible for considerable damage. Khare (1975) pointed out that wilt in lentil, a seed-borne, could be controlled by antagonists, fungicides and irrigation routine management. It also adversely affected that germination of pea seeds (Lambhate and Bhide,1976). Peronospora manshurica was found to be seed-borne on soybean (Rosaca, 1976; Pathak et al., 1978) particularly in seeds

lots imported from the U.S.A. (Mukerwar et al., 1980). Another very important pathogen Glomerella sp. was also reported to be seed-borne on soybean by Roy (1982). Maude and Humpherson-Jone (1977, 1980) observed that seeds of Brassica oleracea were heavily infected with A. brassicicola. Tripathi and Kaushik (1984) while studying the seed mycoflora of rape seed and mustard observed that intensity of infection of A. brassicae seed depended on the lesions on the fruits. Systemic seed transmission of another species of Alternaria i.e. A. sesami in sesame was reported by Dalbir Singh et al. (1983). Presley and Maude (1977a), Maude and Presley (1977) and Bochow (1980) pointed out that neck rot fungus Botrytis allii of onion was internally seed-borne and the infection persisted more than three years. Glushchenko (1980) observed that Peronospora destructor a seed-borne fungus caused epiphytotics of onion in the field. Dhingra (1978) concluded that Fusarium semitectum and Phomopsis spp. were internally seed-borne in bean seeds. Karwarsa and Mohinder (1982) were able to isolate 9 fungi and one bacterium (Xanthomonas cyamopsidis) from certified as well as local variety of cluster beans. Seed-borne nature of Rhizoctonia solani has been reported on cowpea (Diaz-Polanco et al., 1978) and beans (Wanzainum Wan Nik and Yap, 1979). Kamal and Verma while isolating fungi from Pigeon pea seed observed that culture filtrates of some of

the fungi adversely affected the germination of seeds. Out of the three species of Aspergillus isolated from arhar seed by Shukla and Bhargava (1979), A.flavus proved to be pathogenic. Washings of Asparagus seeds were found to contain fungal bodies of Fusarium oxysporum f. sp. asparagi and F.moniliforme (Inglis, 1979).

Damping-off in chillis caused by Colletotrichum capsici has also been reported to be transmitted through seeds (Grover and Bansal, 1971). Colletotrichum sp. and Diaporthe phaseolorum var. sojae have been isolated from soybeans (Girard 1979) and Verticillium alboatrum from lucerne seed (Sheppard and Needham, 1980). Christensen (1983) isolated V.alboatrum in 5 of 20 seed lots of alfa-alfa in Columbia. Srivastava and Gupta (1982) pointed out that seeds of Verbena hybrida collected from ten Indian locations were heavily infested with fungi and some of which were present in embryo and pericarp. Significance of seed-borne pathogens Macrophomina phaseoli, Cercospora cassiicola and Alternaria sesami have been shown by Kushi and Khare (1979). Seed transmission of Macrophomina phaseolina on sesame was reported by Singh and Singh (1982). Out of three species of Fusarium isolated from tomato seeds, F.oxysporum var. herbarum proved to be pathogenic (Orlov et al., 1982) Alternaria alternata f.sp. lycopersici a causal agent of

tomato wilt epidemic was transmitted through seeds (Freshchini, 1982). Seeds obtained from both healthy and blighted tomatoes of 3 field cultivars revealed the presence of Phytophthora infestans (Vartanian & Endo 1985). Inaba et al. (1983) while examining the seed washings to seed mycoflora observed the presence of Oospores of Peronospora farinosa a causal agent of spinach downy mildew in 6 of the eleven cultivars of commercial type studied.

Humpherson-Jones (1943) observed that out of 30 commercial vegetable seed lots tested 13 were found infected with Alternaria brassicae and 15 with Leptosphaeria maculans. Ferreira and Knox-Davies (1984) isolated Rhizopus sp. and Fusarium oxysporum from sweet melon seeds. Srivastava (1984) reported 35 fungi from 40 coriander seed lots. Of these about 17 fungi were pathogenic either to seed or at seedling stage.

The damage to seed by seed-borne pathogens is influenced by several factors. Christensen and Dorworth (1966) pointed out that moisture content of seed play an important role in the establishment of storage fungi in soybean seed. Christensen (1967) also noted an increase in infection by fungi on soybean at 25°C during storage. Similar results were obtained by Kudrina (1967) and Sanchez-Dominguez (1971) while working on seeds of vegetables. Soybean seed samples from different places in Illinois exhibited varied rate of

germination which was attributed to intensity of mycoflora (Tenne et al., 1974). A correlation of photoperiod on resistance of soybean seed with purple seed stain pathogen Cercospora kikuchi in Puerto Rico was noted by Hepperley (1984). Maude et al. (1985) reported that the leaf blight pathogen Alternaria dauci on carrot was favoured by warm, wet conditions and caused considerable damage by defoliation of plants.

Jensen (1967) reported R. carotae as seed-borne and probably this was the first record of this disease on carrot. Several diseases of seed-borne nature on vegetables have been worked out by several workers e.g. Septoria spp. in celery (Hewett, 1968), Botrytis spp. and Fusarium moniliforme var. subglutinosa (Lutynska, 1968 and Vannaci, 1981), Macrophomina phaseolina in cowpea and soybean (Sachston, 1969 and Gangopadhyaya ^{et al.} 1970), Phoma spp. in pigeon pea (Lopezrosa, 1969).

Antibiotics have also been found effective in minimizing the loss due to seed-borne diseases. Sowell (1965) by using streptomycin controlled Xanthomonas sp. on guar while Colletotrichum dematium var. truncata remained unaffected. Rauckyte (1965) could obtain inhibition of Ascochyta fabae on broad bean with nystatin and tetracyclin. Phytobacteriomycin seed treatment and trichothecin dusting on pea and chickpea reduced losses due Fusarium oxysporum and at the same time improved the germination (Kuzmina, 1966). Biomyacin, tetracyclin and terramycin reduced the growth of Pseudomonas lachrymans and Bacillus mesentericus in vitro (Avakyan and Gyusyan, 1966). In addition to above 12 percent biomyacin has been found effective against alternariosis of cabbage (Dorozhkin et al., 1972), griseofulvin and kusumin against verticillium wilt of cotton, griseofulvin and polyoxin against fusariosis of tomato, and griscofulvin against anthracnose of cucumber (Panteleev et al., 1973). Streptomycin against Pseudomonas pisi on pea (Taylor, 1973); Penicillin or streptomycin against P. lachrymans (Kutova and Filipova, 1977); takedamycin and kasugamycin against several fungi on cabbage, carrot, lettuce, radish and spinach, where as bacteria with agrimycin-100 (LO, 1977); an antibiotic obtained from Streptomyces sp. 18 against Pythium debaryanum on tomato, lettuce and cabbage (Numic' et al., 1979), fusamycin and sucamycin against root rot fungi on bean.

Simple treatment of seeds with water at room temperature and hot water treatment have been found to reduce the seed mycoflora (Tyner, 1951). Maude (1966) observed that treatment of pea seeds with steam/air mixture at 65°C or 75°C killed the Ascochyta pisi and A. pinodes. Treatment of pea seeds infected with Ascochyta pinodes with UV radiation improved the germination (Peresypkin, 1966). Hot water treatment has also been found effective against black leg agent Phoma lingam on cabbage (Williams, 1967); against seed-borne cucumber mosaic virus (Sharma and Chohan, 1971); damping-off and stage root rot of broad bean (Elarasi, 1971); Phoma betae and other microbial contaminant on sugar beet (Herr, 1971; Lambat et al., 1974); Fruit rot Rhizoctonia spp. and Choenophora cucurbitarum on okra (Tai Luang Haun and Jamail, 1975); Peronospora parasitica downy mildew and black leg Leptosphaeria maculans on cabbage (Klinkovskaya, 1976), against Macrophomina phaseolina and Fusarium equiseti on cowpea (Sinha and Khare, 1977) Alternaria radicina on carrot (Utyugova and Levteeva, 1977); Fusarium oxysporum f.sp. lagenaria on bottle gourd (Kuniyasu and Nakamura, 1978) and Alternaria brassicae on mustard (Randhawa and Aulakh, 1984). In a modification, the hot water treatment has been combined with fungicide as well treatment which has considerably improved the germination (Wells and Merwarth, 1973; Utyugova and Levteeva, 1977).

Plant products have been reported very effective in minimising the seed mycoflora. Nene et al. (1968) tested antifungal activity of extracts of 88 plants against Helminthosporium turcicum of which 14 exhibited complete inhibition and nine partial inhibition in growth of the fungus. Khandelwal and Prasad (1968,70) used plant extract and fungicides for controlling the seed mycoflora of cucurbits. Kumar and Nene (1969) observed inhibition of Helminthosporium maydis and several other fungi by the extract of Cleome isocandra L. Staron et al. (1969) reported that triterpanic hiteroside (anagalloside) extracted from the stem and leaves of Anagallis arvensis var. phaenicea was inhibitory against Colletotrichum lindemuthianum and Pythium ultimum. Bamode and Shukla (1973) while testing the extracts of 36 plants against 6 pathogens found that pomegranate fruit rind, Plumbago zeylanica root bark adversely affected the growth of several fungi. Singh et al. (1979) could free the gram seeds from wilt fungus Fusarium oxysporum f. sp. Ciceri and Sclerotinia sclerotiorum by treating with aqueous extract of garlic seed.

Culture filtrates could also reduce the seed surface mycoflora. Dimovich et al. (1971) succeeded in reducing the seed infection of pea from Fusarium oxysporum by treating the seed with culture filtrate of 2 species of Penicillium.

Vishunavat and Shukla (1981) reported that culture filtrate of Aspergillus fumigatus increased the germination of lentil and improved the seedling length. Fungi used as antagonists also reduced/controlled the seed borne pathogens in some crops. Akhtar (1969) pointed out that seed decay and damping off caused by Rhizoctonia solani on mustard could be controlled by the action of certain saprophytic soil-borne fungi. Roy and Abney (1979) observed a reduction in incidence of Diaporthe phaseolorum var sojae a seed-borne fungus when inoculated with Cercospora kikuchii. Windles and Kommendahl (1978) used Penicillium oxalicum to control seedling blight of pea. Herman et al (1980) observed reduction in the incidence of Pythium spp. or Rhizoctonia solani on radish and pea with Trichoderma hamatum. Wu and Lu (1984) pointed out that treatment of seeds with Gliocladium virens-19, Trichoderma harzianum-22, T.harzianum-50, Penicillium oxalicum-76 increased emergence and improved seedling health in crucifers and protected them against Alternaria brassicicola. Lifshitz et al (1985) obtained reduction in the incidence of Rhizoctonia pre-emergence damping off of radish by treating seeds with conidial suspension of T.harzianum.

Seed mycoflora can be controlled by several means i.e. physical and chemical means. Mechanical separation of fungal propagules and other debris has been practised for a long time (Anonymous, 1966).

Hot water treatment of seeds has given a good control of seed mycoflora. The basic idea behind heat treatment is that dormant organs such as seeds, bulbs, tubers etc. used as propagative material can withstand higher temperature than those in which their respective pathogen can survive for a given period of time. The temperature of the hot water used and the duration of the treatment varies with different host pathogen combinations. Thus to control loose smut of wheat the seeds are recommended to be treated with hot water for eleven minutes at 58°C (Agrios, 1969). Whereas bulbs are treated at 43°C for 3 hours to get rid off Ditylenchus dipsaci (Agrios, 1969). Low temperature treatment does not kill the pathogen but inhibit or retard the growth and activities thereby prevent the spread of existing infections and initiation of new ones (Agrios, 1969).

Seed treatment with chemicals has been used for centuries for controlling plant pathogens. As early as 1660, salt brine was accidentally discovered to control the bunt of wheat. At the end of 18th century Abb'e Tessier in France tested many chemical compounds to improve wheat seeds,

without affecting agricultural practice (as quoted by Woolmen and Humphrey, 1924). It was Prevost (1807) who demonstrated that copper sulphate could be used against bunt of wheat. It is this study which triggered research for finding out more and more chemicals to control seed-borne pathogens. Formaldehyde was later found to be a good and effective for seed treatment against various seed-borne fungi, particularly smuts (Geuter, 1895; Bolley, 1897). Copper carbonate was found satisfactory as dusting material for controlling wheat smut (Darnell - Smith 1910, 1915-1917). The modern era of fungicides and seed treatment started with the introduction of the organo-mercuric compounds. There has been tremendous amount of literature on this aspect which has been reviewed by Horsfall and Howard, 1959.

It is difficult to summarise all what has been done on chemicals as seed treatment but in the following pages attempt has been made to tabulate some of the information available of chemical seed treatment against seed mycoflora of vegetables.

BENZIMIDAZOLE FUNGICIDES (SYSTEMIC)

Chemical name :- Methyl-N-(1-butylcarbomoyl)-2-benzimidazole carbamate

Trade name :- Benomyl/Benlate

Crop	Disease/pathogen	dosage	References
Pea	<u>Ascochyta disease</u>	42.2-56.6g/ 12.7 kg seed	Maude and Kyle (1970)
"	Powdery mildew	0.3 - 0.5%	Jhooity and Behar (1972)
"	<u>Ascochyta pisi</u>	-	Maude et al (1973)
"	<u>Macrosporaerella pinodes</u>	400 g/kg seed	Yoshii (1975)
"	<u>Fusarium oxysporum</u> f.sp. <u>pisi</u>	-	Lorenzini et al (1975)
"	Root rots and <u>Ascochyta</u> disease	2kg/ton seed	Kotova (1977)
"	Seed mycoflora	-	Batalova and Zino'ev (1978)
"	<u>Fusarium oxysporum</u> f.sp. <u>pisi</u>	-	Utikar et al (1978)
Cucumber	<u>Ascochyta melonis</u> , <u>Colletotrichum lagenarium</u>	0.2 - 0.3%	Oliker (1972)
Sweet melon	<u>Fusarium oxysporum</u>	-	Fereirra and Knox-Davles (1984)
Onion and garlic	<u>Sclerotium cepivorum</u>	10-40g/kg seed	Lafon and Bugaret (1970)
"	<u>S.cepivorum</u>	150g/kg seed	Enwistle and Muna-Singhe (1973)
"	<u>Botrytis allii</u>	1 g/kg	Maude and Presley (1977)

Contd..

Crop	Disease/pathogen	dosage	References
Bean	<u>Rhizoctonia solani</u>	0.1%	Segura and Diazpolanco (1975)
"	<u>Fusarium solani</u> f.sp. <u>phaseoli</u>	-	Russel and Musa (1977)
"	<u>Macrophomina phaseolina</u>	-	Petkar et al (1977)
"	<u>Fusarium</u> spp.	-	Shukla and Bhargava (1978)
Broad bean	<u>R.solani</u> and <u>F.oxysporum</u>	4g / kg seed	Eisa and Brakat (1978)
Faba bean	<u>Ascochyta faba</u>	0.2%	Bernier (1979)
French bean	<u>Phaeoisariopsis griseola</u>	-	Sindhan (1984)
Bean	Chocolate spot	-	Bainbridge et al (1985)
Chickpea	<u>Ascochyta rabeiei</u>	-	Kaiser et al (1973)
"	<u>A. rabeiei</u>	-	Reddy (1980)
"	<u>Fusarium oxysporum</u> f.sp. <u>ciceri</u> and <u>F.solani</u>	-	Mani and Sethi (1984)
Pea	<u>Fusarium oxysporum</u> f. sp. <u>pisi</u>	-	Utikar et al (1978)
"	Seed mycoflora	-	Batalova and Zinov'ev (1978)
"	<u>F.oxysporum</u> f. sp. <u>pisi</u>	-	Shukla et al (1979)
Gram	<u>Sclerotium rolfsii</u> , <u>Rhizoctonia bataticola</u> and <u>F.oxysporum</u> f.sp. <u>ciceri</u>	-	Verma and Vyas (1977)
Tomato	Seed mycoflora	-	Nedumaran and Vidyasekharan (1981)

Chemical name :- 2-(4'-thiazolyl)-benzimidazole

Trade name :- Thiabendazole

Crop	Disease/pathogen	dosage	References
Cabbage	<u>Phoma lingam</u>	1,1.2 and 5 a.i/ kg seed	Maude <u>et al</u> (1973)
Onion	<u>Botrytis cinerea</u>	-	Presley and Maude (1980)
Bean	<u>Fusarium solani</u> f. sp. <u>phaseoli</u>	-	Russel and Musa (1977)
Chickpea	<u>Alternaria rabeiei</u>	-	Kaiser <u>et al</u> (1973)

OXATHIIN FUNGICIDES (SYSTEMIC)

Chemical name :- 5,6-dihydro-2-methyl-1,4-Oxathiin-3-carboxinilide

Trade name :- Vitavax/Carboxin

Crop	Disease/pathogen	dosage	References
Cucubrit	<u>Rhizoctonia solani</u>	-	Jhooty and Grover (1971)
Onion	<u>Urocystis magica</u>	25 g/kg	Crete and Tartier (1973)
Chillies	<u>Colletotrichum capsici</u>	-	Narain and Panigrahi (1972)
Bean	<u>Macrophomina phaseolina</u>	-	Petkar et al (1977)
"	<u>Fusarium</u> spp.	-	Shukla and Bhargava (1978)
French bean	<u>Phaeoisariopsis griseola</u>	-	Sindhan (1984)
Carrot	<u>Alternaria dauci</u> , <u>Stemphyllum radicum</u> and <u>S.radicinum</u> var. <u>petroselina</u>	200g/100 kg seed	Nikova (1979)
Gram	<u>Sclerotium rolfsii</u> , <u>Rhizoctonia batati-</u> <u>cola</u> and <u>F.oxysporum</u> f.sp. <u>ciceri</u>	-	Verma and Vyas (1977)
Chemical name :- 2,3-dihydro-5-carboxanilido-6-methyl-1, 4-oxathiin-4, 4-dioxide			
Trade Name :- Plantavax/Oxycarboxin			
Chillies	<u>Colletotrichum capsici</u>	-	Narain and Panigrahi (1972)
Bean	<u>Uromyces appendiculatus</u>	-	Okioga and Joffer (1972)

ORGANIC MERCURIAL FUNGICIDES

Chemical name :- Ethylmercury/Phenylmercury acetate etc.

Trade name :- Cerasan Red/Cereson dry etc.

Crop	Disease/Pathogen	dosage	References
Cabbage	<u>Pythium aphanidermatum</u> , <u>Rhizoctonia solani</u> and <u>R. bataticola</u>	-	Grewal and Singh (1965)
Sugarbeet	<u>P. aphanidermatum</u>	2.5g/kg seed	Sen et al (1974)
Okra	<u>Rhizoctonia bataticola</u>	0.3 %	Vir and Gaur (1970)
"	<u>R. bataticola</u>	0.3 %	Goel and Mehrotra (1973)
Bottlegourd	Germinability of seed	-	Sohi (1976)
Bean	Glomerella glycines	-	Ahn and Chung (1970)
"	Seed mycoflora	-	Kaul (1973)
"	Macrophomina phaseolina	-	Petkar et al (1977)
Faba bean	<u>Fusarium</u> spp.	-	Shukla and Bhargava (1978)

Crop	Disease/Pathogen	dosage	References
	Chemical name :- Phenylmercury acetata + Ethylmercury chloride		
	Trade name :- Agrosan - GN		
Bean	Seed mycoflora	-	Kaul (1973)
"	<u>M. phaseolina</u>	-	Petkar <u>et al</u> (1977)
Faba bean	<u>Fusarium</u> spp.	-	Shukla and Bhargava (1978)
French bean	<u>Phaeoisariopsis griseola</u>	-	Sindhan (1984)
	Chemical name :- Ethyl Mercury chloride		
	Trade name :- Granosan/ 2% cerasan		
Pea and broad bean	Seed mycoflora	2-4kg/ton seed	Volkova (1964)
Cucurbit	<u>Bacillus mesentericus</u> and <u>F. oxysporum</u> var. <u>niveum</u>	3g/kg seed	Avakyan (1966)
Beet	Black leg disease	4kg/ton seed	Pozhar (1971)

ORGANIC SULPHUR COMPOUNDS

Chemical name :- Tetramethyl thiuram disulphide or bis disulphide

Trade Name :- Thiram

Crop	Disease/Pathogen	dosage	References
Pea and Broad beans	Seed mycoflora	2kg/ton seed	Vclkova (1964)
Pea	<u>Ascochyta pisi</u> , <u>Helminthosporum sativum</u> <u>Sclerotinia fructigena</u> & <u>Puccinia triticina</u>	-	Ksendzova (1974)
"	Post and pre-emergence mortality		Bedi & Kapur (1974)
"	Root rots and Ascochyta disease	50%	Kotova (1977)
"	Germination and emergence of plant	100-400g/kg seed	Rivera (1982)
"	<u>Pythium ultimum</u>	-	Lewis and Lumsden (1984)
Cabbage	11 seed-borne pathogens	-	Mande et al (1969)
Brassicas	<u>Pythium</u> spp., <u>Peronospora parasitica</u>	-	White et al (1984)
Cucurbits	<u>Bacillus mesentericus</u> and <u>Fusarium oxysporum</u> var. <u>niveum</u> .	2%, 8-9g/kg seed	Avakyan (1966)
"	Germination of seeds	300-500g/kg	Alkamper (1966)
Beet	Black leg disease	6kg/ton seed	Pozhar (1971)
Onion	Collar rot	-	Boguslawski (1967)
"	<u>Botrytis</u> spp.	-	Lutynska (1968)

Crop	Disease/Pathogen	dosage	References
Onion	<u>Urocystis cepula</u>	-	Macias (1970)
"	<u>Urocystis magica</u>	25g/kg seed	Crete and Tartier (1973)
Chillies	<u>Colletotrichum capsici</u>	0.2%	Chakravarti and Anil Kumar (1975)
"	Some important diseases	-	Aleksic' and Aleksic'(1976)
"	<u>Colletotrichum dematium</u>	0.2%	Siddiqui et al (1977)
"	Pre-and post emergence losses	0.3%	Jharia et al (1978)
Okra	<u>Rhizoctonia bataticola</u>	-	Goel and Mehrotra (1973)
Watermelon & Melon	Improved germination	-	Sohi (1976)
Vegetables	Bacteria and fungi on seed	-	Kolev et al (1976)
Bean	Seed mycoflora	-	Kaul (1973)
"	<u>R.solani</u> and <u>F.oxysporum</u>	4g/kg seed	Eisa and Brakat (1978)
Faba bean	<u>Ascochyta faba</u>	0.2%	Bernier (1979)
Carrot	<u>Alternaria radicina</u> and <u>A.dauci</u>	-	Soteros (1979)
Brinjal	<u>Helminthosporium</u> sp.	-	Sarode and Kadam (1977)
Beet	Damping-off	-	V'rbanov et al (1984)
Onion	Microflora of seed	-	Kaul (1972)
Coriander	Mycoflora of seed	0.3%	Srivastava (1984)

Contd....

Crop	Disease/Pathogen	dosage	References
Gram	Seedling rot	-	Shrisat and Kale (1979)
	Chemical name :- Zinc ethylene bisdithiocarbamate		
	Trade name :- Zineb		
Onion	Collar rot	-	Boguslawski (1967)
Chillies	Some important disease	-	Aleksic' and Aleksic'(1976)
Carrot	<u>Sclerotinia</u> , <u>Botrytis</u> and <u>Rhizopus</u> sp.	0.1%	Tasca and Hulea (1973)
	Chemical name :- Zinc dimethyl dithiocarbamate		
	Trade name :- Ziram		
Chillies	<u>Colletotrichum capsici</u>	-	Narain and Panigrahi (1972)
Okra	Fusarium wilt	0.3%	Kapoor (1978)
French bean	<u>Phaeoisariopsis griseola</u>	-	Sindhan (1984)
	Chemical name :- Manges ethylene bisdithiocarbamate		
	Trade name :- Maneb/Mancozeb/Dithane-22		
Cabbage	Soft rot and curd disease	-	Skowronski (1976)
Beet	<u>Phoma betae</u> and <u>Pythium ultimum</u>	80%	Dorpoux et al (1966)
Gram	Seedling death	-	Gurdip Singh and Bedi (1980)
	Chemical name :- Ferric dimethyl dithiocarbamate		
	Trade name :- Ferbam/Hexaferb		
Onion	<u>Botrytis</u> spp.	-	Lutynska (1968)
Bean	<u>Fusarium</u> spp.	-	Shukla and Bhargava (1978)

HETEROCYCLIC NITROGENOUS FUNGICIDES

Chemical name :- N-trichloromethyl thio-4-cyclohexene-1,2 dicarboxide

Trade name :- Captan/Orthocide

Crop	Disease/Pathogen	dosage	References
Garden pea	bacterial and fungal pathogens	-	Boyadzhiev (1972)
Pea	<u>Fusarium</u> sp.	-	Kirik and Steblyuk (1972)
"	Post and pre-emergence mortality	-	Bedi and Kapur (1974)
"	<u>Fusarium oxysporum</u> f.sp. <u>pisi</u>	0.2%	Gangopadhyay and Kapoor (1976)
"	Pre-emergence damping-off by <u>Pythium ultimum</u>	-	Ohh et al (1978)
"	<u>Ascochyta</u> sp.	-	Kuhne (1983)
Cabbage	<u>Pythium aphanidermatum</u> , <u>Rhizoctonia solani</u> and <u>R.bataticola</u>	-	Grewal and Singh (1965)
Chillies	<u>Colletotrichum capsici</u>	-	Chakravarti and Anil Kumar (1975)
"	Some important diseases	-	Aleksic' and Aleksic'(1976)
"	<u>Phytophthora capsici</u>	-	Aleksic' et al (1976)
"	improvement in germination capacity	-	Sohi (1976)
Bean	<u>Glomerella glycines</u>	-	Ahn and Chung (1970)
"	<u>Colletotrichum lindemuthianum</u>	200g/100kg seed	Petrov (1972)
"	<u>Macrophomina phaseolina</u>	-	Petkar et al (1977)

Contd....

Crop	Disease/Pathogen	dosage	References
Carrot	Seedling emergence	0.3% W/W	Perry and Hegarty (1971)
Sugarbeet	<u>Phoma betae</u> , <u>F. oxysporum</u> and <u>Penicillium</u> sp.	600/100kg seed	V'rbanov (1973)
"	Black leg disease	600g/100 kg seed	V'rbanov (1974)
Beet	Damping-off	-	V'rbanov (1984)
Chickpea	<u>Ascochyta blight</u>	0.1%	Bhatti et al (1983)
"	<u>A. rabeiei</u>	0.1%	Bhatti et al (1984)
Pea	Chemical name :- <u>Cis-N-(1,1,2,2-tetrachloroethylthio)-4-cyclohexene-1,2-dicarboximide</u> Trade name :- Captafol/difolatan	0.1%	Gangopadhyaya and Kapoor (1976)
Tomato	<u>Fusarium oxysporum</u> f.sp. <u>pisi</u>	-	Nedumaran and Vidyasekharan (198
Beet	Seed mycoflora	80%	Dorpoux et al (1966)
Chillies	<u>Pythium ultimum</u>	-	Chakravarti and Anilkumar (1975)
"	<u>Colletotrichum capsici</u>	-	Aleksic' et al (1976)
"	<u>Phytophthora capsici</u>	0.25%	Siddiqui et al (1977)
Okra	<u>Colletotrichum dematium</u>	-	Chauhan et al (1979)
	<u>Rhizoctonia</u> root rot		

MISCELLANEOUS FUNGICIDES

Chemical name :- Pentachloronitrobenzene (PCNB)

Trade name :- Quintozene/Brassiccol

Crop	Disease/Pathogen	dosage	References
Cucurbits	<u>Rhizoctonia solani</u>	-	Jhooty and Grover (1971)
Sweet melon	<u>Rhizopus</u> sp.	-	Fereirra and Knox-Davies (1984)
Sugar beet	<u>Sclerotium rolfsii</u> , <u>Rhizoctonia solani</u>	5g/kg seed	Sen et al (1974)
Beet	<u>R. solani</u>	-	Natti (1966)
Broad bean	<u>R. solani</u> and <u>Fusarium oxysporum</u>	4g/kg seed	Eisa and Brakat (1978)
Carrot	<u>Sclerotinia</u> , <u>Botrytis</u> and <u>Rhizopus</u> sp.	0.1%	Tasca and Hulea (1973)
Onion	<u>Microflor</u> of seed	-	Kaul (1972)
Okra	<u>Rhizoctonia bataticola</u>	-	Goel and Mehrotra (1973)
Gram	Seedling emergence	-	Shirsat and Kale (1979)

Chemical name :- 2,6-dichloro-4-nitroaniline

Trade name :- Bicloran/Botran

Onion	<u>Sclerotium cepivorum</u>	30g a.i./kg seed	Lafon and Bugaret (1968)
"	<u>S. cepivorum</u>	150-300g/kg seed	Enwistle and Muna Singhe (1973)

=====

M A T E R I A L S A N D M E T H O D S

=====

MATERIALS AND METHODS

Seed samples of cauliflower and pea will be collected from the farmers field in and around Aligarh. Seeds will also be procured from National Seed Corporation and Indian Agriculture Research Institute, New Delhi. Seeds so obtained will be stored in air tight containers of P.V.C. Jars.. Initially the seeds will be subjected to visual examination and examination under the microscope after soaking in water for 12 hours, for the presence of contaminants (Conidia, pycnidia and mycelial fragments) on the surface.

Seeds will be transferred in water contained in 100 ml flask and shaken vigorously. The washing so obtained will be centrifuged to obtain the spore/fungal mass. The clear supernatant will be discarded and the residue will be examined microscopically for the presence of the fruiting body.

External seed mycoflora will be studied by using Blotter and Agar plate methods.

1. Blotter Methods :-

Moistened sterilized blotting papers will be placed in sterilized Petridishes of 9 cm. diameter. In each Petriplates equal number of seeds will be placed equidistantly

The petridishes will be incubated at $20 \pm 2^{\circ}\text{C}$ for 8 days with cycles of 12 hours light and 12 hours darkness. After 8 days of incubation fungi developing on the seeds will be isolated and identified.

2. Agar Plate Method:-

The seeds will be transferred in Petridishes of 9 cm diameter containing plain agar medium at equal distance. Incubation will be done as in the case of Blotter Method.

For internal seeds mycoflora both Blotter and Agar Plate Methods will be used. The seeds will be surface sterilized in 1:1000 mercuric chloride for 2-5 minutes followed by washings with sterilized water and will be transferred to Petriplates. Throughout the studies the plates will be incubated at $20 \pm 2^{\circ}\text{C}$. During the course of examination, the percentage germination of seeds and frequency and relative abundance of fungi will be determined. In these studies more attention will be paid to the mycoflora of structurally abnormal seeds. In internally seed-borne mycoflora, the site of infection in the seeds will be determined. Surface sterilized seeds with 1:1000 mercuric chloride will be placed in Petridishes containing plain agar and incubated at $20 \pm 2^{\circ}\text{C}$. The seeds showing mycelial infection will be boiled in KOH for 5 - 10 minutes till

they appear soft and translucent. The seeds will be given several washing with lactophenol and will be macerated and examined under the microscope.

In order to determine the pathogenicity of fungi appearing in these studies the seeds (100 in number) after having surface sterilized will be inoculated by placing them over the culture of potential pathogen grown on PDA and will be transferred to Petriplates. Equal number of inoculated seeds will also be kept to serve as control. Germination percentage and the development of symptoms if any will be observed during the course of these studies. In case fungi show some potential for pathogenesis the pathogenicity will also be determined at seedling stage. The seedlings will be inoculated by pouring suspension of the 5 gram fungus into the soil.

Effect of culture filtrates of pathogenic fungi on germinability of seeds will also be tested. The fungus will be grown in Richards medium contained in 250 ml. flask for 15 days. Surface sterilized seeds (100 in number) will be soaked in fungal culture filtrate for 24 hours. After soaking, seeds will be washed with sterilized distilled water and placed in sterilized Petriplates containing moist blotter paper. Appropriate number of seeds will be treated with sterilized distilled water to serve as control. Number of seeds germinated will be recorded as to calculate the percentage germination.

Attempts will be made to control the seed mycoflora with fungicides, nematocides, antibiotics and extract of certain plants known for antimicrobial activity. The seeds will be treated with fungicides, antibiotics and plant extract at different percentage of dilution and different time intervals and will be plated in sterilized petriplates. The plates will be incubated at $20 \pm 2^{\circ}\text{C}$ for a week. The plates will be examined regularly for germination of the seed and the development of any fungi. Untreated seed will also be kept for control. Frequency and relative abundance of fungi will be calculated.

To test the effect of neem leaves on storage fungi, fresh neem leaves obtained will be oven dried at 60°C and will be cooled at room temperature. The leaves will be powdered and mixed with the seeds to be stored.

Through out the studies a lot of 400 seeds per sample will be taken unless stated otherwise. The data so obtained will be analysed statistically. The frequency and relative abundance of fungi will be calculated as under.

$$\text{Frequency} = \frac{\text{No. of seeds containing a particular fungus}}{\text{Total No. of seeds used}} \times 100$$

$$\text{Relative abundance} = \frac{\text{Total no. of colonies of a fungus}}{\text{Total no. of colonies of all the fungi}} \times 100$$

PLAN OF WORK

It is proposed to study the following aspects.

1. Survey of the seed-borne diseases of some important vegetable crops viz. cauliflower and peas.
2. Study of the external and internal mycoflora of cauliflower and peas.
3. Location of pathogens in the seed and their role in transmission of the diseases.
4. Pathogenicity studies of selected pathogenic fungi isolated from seeds of cauliflower and peas.
5. Effect of culture filtrate of important pathogenic fungi on germination of seeds of cauliflower and peas.
6. Effect of different parts and products of various plants known for antimicrobial activity on the seed mycoflora.
7. Effect of the above with fungicides on the seed-mycoflora.
8. Effect of dried leaves of neem (Azadirachta indica Juss.) on seed mycoflora.

=====

R E F E R E N C E S

=====

REFERENCES

- Abdalla, M.H.(1974). Mycoflora of ground nut Kernal s from the Sudan. Trans. Brit. Mycol. Soc. 63(2): 353-359.
- Agrios, George N.(1969). "Plant Pathology". Academic Press New York and London.
- Ahn, J.K. & Chung, B.K.(1970). Effects of seed disinfectants for controlling the soybean anthracnos. Jnl. Pl. Prot. Korea, 9(1): 21-24.
- Akhtar, C.M.(1969). Biological control of Rhizoctonia solani causing seed decay and damping off of Mustard. W.Pakist. Jn. Agric. Res. 7(4): 29-52.
- Aleksic, Z., Aleksic, D. and Sutic, D.(1976). The possibility of control of Phytophthora capsici Leonian, the pathogen of pepper blight. Zastita Bilja, 27(137/138): 257-266.
- Aleksic, Z. and Aleksic, D.(1976). The control of the most important disease of pepper. Poljoprivreda, 23(252): 24-31.
- Alkamper, J.(1966). The influence of seed dressing on germination and sprouting capacity in pumpkin. Nachr. Bl. dt. pflschutzdienst., Stutlf., 18(7): 105-109.
- Anderson, T.R.(1985). Seed moulds of soybean in Ontario and the influence of production area on the incidence of Diaporthe phaseolorum var. caulivora and Phomopsis sp. Canad. Jour. Plant Path., 7(1): 74-78.
- Anonymous (1941). Fusarium root rot of beans. Agric. Gaz., New South Wales 52: 41-42.

- Anonymous (1965). Losses in agriculture. Agric. Res. Serv., Agric Handbook No.291, Washington. 120p.
- Anonymous, (1966). International Rules for Seed Testing. Proc. Int. Seed Test. Ass. 31: 1-152. -
- Askerov, I.B.(1968). Askokhitoz Nuta (Ascochyta of Chickpea) Zashch. Rast. Moscow 13: 52-53.
- Avakyan, S.A.(1966). Influence of some seed preparations on the causal agent of cucurbita disease. Erevan, 3(13): 220-230.
- Avakyan, S.A. & Gyusyan, M.M.(1966). On the influence of some antibiotics on the causal agent of disease of cucurbits. Erevan, 3(13): 231-238.
- Bainbridge, A., Fitt, B.D.L., Creighton, N.F. and Cayley, G.R. (1985). Use of fungicides to control chocolate spot (Botrytis fabae) on winter field beans (Vicia faba) Plant Pathology, 34(1): 5-10.
- Bambode, R.S. and Shukla, V.N.(1973). Antifungal properties of certain plant extracts against some fungi. Punjabaro Krishi Vidyapeeth Res. Jour., 2(1): 1-8.
- Barros, S.T.DE, Menenzes, M., Fernandes, M.J. and Lira, N.P. (1985). Seed-borne fungi of 34 Cowpea (Vigna unguiculata) cultivars in the state of Pernambuco, Brazil. Fitopatologia Brassileira, 10(1): 85-95.
- Bassi, A.JR., Goode, M.J. and Dowers, J.L.(1978). Spinach Fusarium decline, a wide spread and destructive disease in Arkansas. Arkansas Farm. Research, 57(2): 13.
- Batalova, T.S. and Zinov'ev, L.S.(1978). Trials of fungicides seed protectants safe for nodule bacteria. Byulleten' Vsesoyuznogo Nauchnogo Nauchno-Issledovatel'skogo Institute Zashchity Rastenii, No.43: 28-32.

- Bedi, P.S. and Kapur, K.S.(1974). Mycoflora of pea and its associative effects on the crop. Jour. Research, 11(3): 284-288.
- Bedlan, G.(1985). Pea leaf and Pod spot. Pflanzenschutz, No. 147, 14-15.
- Bernier, C.C.(1980). Fungicidal control of Ascochyta blight of faba beans. Faba Bean Information Service Newsletter No.2, 43.
- Bhatti, M.A., Hussain, S.A. and Malik, M.(1984). Effect of fungicidal treatment of diseased seed against Ascochyta rabiei on germination in chickpea. Pak. Jour. Agric. Res. 5(1): 23-25.
- Bhatti, M.A., Malik, B.A. and Hussain, S.A.(1983). Efficacy of Calixin-M and other fungicides against Aschchyta blight in chickpea. International Chickpea Newsletter No.8, 23-24.
- Bochow, H.(1981). Investigation of seed-borne transmission of Botrytis allii Munn in Onion (Allium cepa) and the influence of seed dressing with fungicides. Archiv fiir Phytopathologe und Pflanzenschutz 17(1): 31-38.
- Boguslawski, W.(1967). Preliminary investigation on the efficacy of treatment of Onion seed bulbs against collar rot (B.allii). Pr. nauk. Inst. Ochr. Rosl., 9(1): 195-198.
- Bolley, H.L.(1897). New work upon smut of wheat, oats and barley and a resume of treatment experiments for the last three years. North Dakota Agric. Exp. Sta. Bull. 27: 109-164.

- Boyadzhiev, KH.(1972). Seed treatment of Garden bean with fungicides and antibiotics. Rastitelna Zashchita, 20(8): 26-30.
- Carter, W.(1962). Insects in Relation to Plant Diseases. Inter Science Publishers, New York and London.
- Chakravarti, B.P. and Anil Kumar, T.B.(1975). Control of seed-borne infection of Colletotrichum capsici in chillies. Current Research, 4(10): 172.
- Chang, H.C.(1960). Primenenie fungitsidovi antibiotikov v bor'be s abtraknozom Fasoli. (The use of fungicides and Antibiotics for the control of anthracnose of bean) Zashchita rastenii ot vreditel'ei i boleznei. Mem. Inst. agron. Leningr. 80: 160-163.
- Chattopadhyay, S.B. & Mustafee, T.P.(1967). Outbreaks and new records, India. Occurance of two new diseases on vegetable. Pl. Prot. Bull., F.A.O., 15(2): 40-41.
- Chauhan, M.S., Tripathi, N.N., Duhan, J.C., Virk, K.S.(1979). Rhizoctonia root rot of okra and its control. Pesticides, 13(1): 24-25.
- Cherewick, W.J.(1954). Studies on seed-borne microflora and the effect of seed treatment of rice. Malay, Agric. 27(3): 169-172.
- Christensen, A.A.(1983). Incidence of external seed-borne Verticillium albo-atrum in commercial seed lots of alfa-alfa. Plant Disease 67(1): 17-18.
- Christensen, C.M.(1951). Fungi on and in wheat seed. Cereal. Chem. 28(5): 408-415.
- Christensen, C.M.(1967). Increase in invasion by storage fungi and in fat acidity value of commercial lots of soybeans stored at moisture content of 13.0% - 14.0% Phytopath., 57(6): 622-624.

- Christensen, C.M. & Dorworth, C.E.(1966). Influence of moisture content, temp. and time of invasion in soybean by storage fungi. Phytopath. 56(4): 412-418.
- Cramer, H.H.(1967). Plant Protection and World Crop Production Leverkusen: Bayer Pflanzenschutz. 524p..
- Crete, R. and Tartier, L.(1973). Three years of chemical control trials against smut of Onion, Urocystis magica Phytoprotection, 54(1): 31-40.
- Crosier, W.F.(1961). The paradoxical value of seed-borne fungi Ass. Off. Seed. Anal. Newsletter, 35: 14-17.
- Dalbir Singh, Mathur, S.B. and Neergaard, P.(1983). Systemic seed transmission of Alternaria sesamicola in Sesamum indicum. Trans. Brit. Mycol. Soc. 80(3): 570-571.
- Darmell-Smith, G.F.(1917). The prevention of bunt, experiments with various fungicides. Agric. Gaz. New South Wales, 28: 185-189.
- Darpohz, H., Lebrun, A. and De La Tullaye, B.(1966). Efficacy of fungicides in Beet seed disinfection, I.I.R.B., 1(2): 67-83.
- De Tempe, J.(1964a). Seed-borne Fusarium infection in temperate climate cereals. Proc. Inst. Seed Test. Ass. 29: 117-140.
- De Tempe, J.(1964b). Helminthosporium sp. in seed of wheat, barley, oats and rye Proc. Int. Seed Test. Ass. 29: 117-140.
- D'eroCole, N. and Sportelli, M.(1982). Mycoflora on Cicer arietinum seeds. Informatore Fitopatologico, 32(6): 51-54.

- Dhingra, O.D., Nichololson, J.F. and Sinclair, J.B. (1973).
Influence of temperature on recovery of Aspergillus flavus from soybean seed, Pl. Dis. Reprtr. 57(2): 185-187.
- Dhingra, O.D. (1978). Internally seed-borne Fusarium semitectum and Phomopsis sp. affecting dry and snap bean seed quality. Pl. Dis. Reprtr. 62(6): 509-512.
- Diazpolanco, C., Maurezutt, P. and Salasdeziaz, (1978).
Rhizoctonia solani, pathogen on cowpea (Vigna unguiculata) in Venezuela. Agronomia tropical, 28(4): 409-418.
- Dimovich, V.O., Kirik, M.M. and Steblyluk, M.I. (1971). Effect of antibiotics of some penicillium strains on the growth of pea and its susceptibility to Fusarium disease. Mikrobiologichnii Zhurnal, 33(5): 558-561.
- Dorozhkin, N.A., Kustova, A.I. and Nitievskaya, V.I. (1972).
The effect of Trichoderma, antibiotics and phytoncides on infection of cabbage by diseases. Kul'tur, No.2: 96-101.
- Dwivedi, R.S. and Tandon, R.N. (1976). Studies on mycoflora of stored seeds of bottle gourd. Acta Betanica Indica 4(2): 139-143.
- Eisa, N.A. and Brakat, F.M. (1978). Relative efficiency of fungicides in the control of damping-off and stemphylium leaf spot of broad bean (Vicia faba). Pl. Dis. Reprtr. 62(2): 114-118.
- Elarosi, H., Abd-El-Rehim, M.A., Michail, S.H. and El-Ahmad, A.M. (1971). Hot water treatment as a method for decreasing damping-off, and early stages of root rot of Horse Bean. Phytopath. Mediterranea, 10(1): 94-106.

Ellis, M.A.; Galvez, G.E. and Sinclair, J.B.(1976). Internally seed-borne fungi and seed quality of bean (Phaseolus vulgaris L.) harvested in small farms in four department of columbia. Noticias Phytopatologicas, 5(2):79-82.

Enwistel, A.R. and Muna Singhe, H.L.(1973). 23rd Annual Report for 1972, National Vegetable Research Station Wellesbourne, Warwick.

Fereirra, J.F. and Knox-Davles, P.S. (1984). Occurrence and control of Fusarium oxysporum on sweet melon seed. Phytophylactica, 16(1): 67-69.

Frank, B.(1883). Ueber einige neue und weniger bekannte pflanzenkrankheiten. Ber. deutsch. Bot. Cessell. 1: 29-34.

Frenschini, A., Corda, P. and Cartar, C.(1982). Observation on the pathogenesis and epidemiology of tomato (Lycopersicum esculentum Mill.) wilt due to Alternaria alternata f.sp. Lycopersici Grogan, Kimble & Misaghi Studi Sassresi III, 29: 61-66.

Gambogi, P.(1964). On the mycoflora of beet seed clusters. Annali Sper.Agr. N.S. 18(5-6): 815-849.

Gangopadhyaya, S. and Kapoor, K.S.(1976). Wet seed treatment for control Fusarium wilt of pea. Vegetable Science, 3(1): 74-78.

Gangopadhyay, S., Wyllie, T.D. and Luedders, V.D.(1970). Charcoal rot disease of soybean transmitted by seeds. Pl. Dis. Reprtr. 54(10): 1088-1091.

Geuter, Th.(1895). Uber die Finwirkung van Formaldehydelosugen auf Getreidebrand. Ber.Pharm. Gesell. 5: 325-329.

- Gibson, I.A. (1957). Saprophytic fungi as destroyers of germinating pine seeds. F. Afr. Agric. J. 22:203-206.
- Giessmann, H.J. and Daebler, F. (1973). Studies on the biology and control of the fungus Phoma lingam (Tode ex Fr.) Desm on cabbage. Archiv. fur, Phytopathologie und Pflanzenschutz 9(1): 5-13.
- Girard, J.C. (1979). Detection of two diseases on soybean seeds from senegal. Agronomie Tropicale, 43(3): 305-307.
- Glushchenko, V.I. (1980). The seed transmission downy mildew of onion. Zashchita Rastenii No.8: 31.
- Goel, S.K. and Mehrotra, R.S. (1973). Rhizoctonia root rot and damping-off of okra and its control. Acta Botanica Indica, 1($\frac{1}{2}$): 45-48.
- Goel, S.K. and Mehrotra, R.S. (1977 publ. 1978). Root and collar rot of okra caused by Rhizoctonia bataticola and its control. Indian Phytopath., 30(1): 112-113.
- Gomes, J.L.L. and Dhingra, O.D. (1983). Alternaria alternata a serious pathogen of white colored snap bean (Phaseolus vulgaris) seeds. Fitopatologia Brasileira 8(1): 173-177.
- Grewal, J.S. and Singh, R.P. (1965). Chemical treatment of seed and nursery bed to control damping-off of cabbage. Indian Phytopath., 18(3): 225-228.
- Grover, R.K. and Bansal, R.D. (1971). Seed-borne nature of Colletotrichum capsici in chillies seeds and its control by seed dressings fungicide. Indian Phytopath. 23(4): 664-668.

- Gupta, V.K. and Chohan, J.S.(1971). Losses and nature of damage caused by seed rot fungi in stored Groundnut in Punjab. Seed-borne fungi and seed health testing in relation to seedlings disease of groundnut. Indian Phytopath. 23(4): 603-605, 622-625.
- Gupta, V.K. and Saharan, G.S.(1973). Seed rot and root rot complex of beans (Phaseolus vulgaris L.) Biologica Plantarum 15(2): 123-125.
- Gurdip Singh and Bedi, S.P.(1979 publ. 1980). Control of foot rot of gram caused by Operculella padwickii, Indian Phytopath. 32(3): 491-493.
- Halfon-Meiri, A.(1970). Infection of chickpea seeds by Ascochyta rabiei in Israel. Pl. Dis. Reprtr. 54: 442-445.
- Harman, G.E., Chet, I. and Baker, R.(1980). Trichoderma hamatum effects on seed & seedling disease induced in radish and pea by Pythium sp. or Rhizoctonia solani Phytopathology, 70(12): 1167-1172.
- Harrison, J.G.(1978). Role of seed-borne infection in epidemiology of Botrytis fabae on field beans. Trans. Brit. Mycol. Soc. 70(1): 35-40.
- Haware, M.P.(1971). Fungal microflora of seeds of Pisum sativum L. and its control. Mycopath. Mycol. appl. 43(3-4): 343-345.
- Hellwig, L.C.(1699). Kurzes send Schreiben Wegendes Sogenaten Honigthaues Welcher Sick an hewigen Korn Sehen-Lassen, Und die grosen Schwarzen Korner inegemein Mutter Korn gennent, hervogebraucht, was de van Zu hatten, Woven as entstanden und abes niitzlichoder schadlich say (Langensalze, 1699, Cited by C.Berger, Ergot and Ergotin, 1931).

- Hepperley, P.R.(1984). Purple seed stain incidence among soybean cultivars and between seasons in Puerto Rico. Jour. Agric. of the University of Puerto Rico 18(1): 87-99.
- Hepperley, P.R. and Rudrigues, R.(1984). Mycofloral succession and viability losses in pigeon pea seed in Puerto Rico. Jour. Agric. of the University of Puerto Rico 68(1): 19-31.
- Herr, L.J.(1971). Hot water treatment for elimination of seed borne Phoma betae and other microbial cantaminants from sugar beet seed. Jnl. American Soc. of Sugar Beet Technologists, 16(7): 568-574.
- Hewett, P.D.(1964). Carrot seed and two fungus diseases. Agriculture, London, 71(12): 555-556.
- Hewett, P.D.(1966). Ascochyta fabae spg. on Tick Bean seed. Plant Pathology 15(4): 161-163.
- Hewett, P.D.(1968). Viable septoria spp. in celery seed samples. Ann. appl. Biol. 61(1): 89-98.
- Hewett, P.D.(1973). The field behaviour of Ascochyta fabae, and Disease control in field beans. Ann. appl. Biol. 74(3): 287-295.
- Horsfall, J.G. and Howard, F.L.(1959). Therapy in "Plant Pathology" (J.G.Horsfall and A.E.Diamond eds.) vol.I, PP. 563-604.
- Holtzhausen, M.A. and Knox-Davles (1974). Pathogens of cruciferous vegetable crops in commercial seed samples in South Africa. Phytophylactica, 6(4): 289-294.

- Holtzhausen, M.A.(1978). Seed-borne fungal pathogens and diseases of Japanese radish and their control in South Africa. Phytophylactica, 10(4): 107-113.
- Humpherson-Jones, F.M.(1983). Pathogenicity studies on isolates of Leptosphaeria maculans from brassice seed production crops in South-east England. Ann. appl. Biol. 103(1): 37-44.
- Inaba, T., Takahashi, K. and Morinaka, T.(1983). Seed transmission of Spinach downy mildew Plant Disease, 67(10): 1139-1141.
- Inglis, D.A.(1979). Contamination of asparagus seed by Fusarium oxysporum f.sp. asparagi and Fusarium moniliforme. Plant Disease, 64(1): 74-76.
- Jain, J.P. and Patel, P.N.(1969). Seed mycoflora of Gaur, their role in emergence and vigour of seedlings and efficacy of fungicides. Indian Phytopath. 22(2): 245-250.
- Jensen, A.(1967). A new and important disease in Danish carrot colt storage. Ugeskr. Agron., 112(35): 659-665.
- Jharia, H.K., Khare, M.N. and Chand, A.(1977 publ. 1978). Indian Phytopath., 30(3): 341-343.
- Jhoopty, J.S. and Behar, D.S.(1972). Control of Powdery mildew of pea and pumpkin by seed treatment with benomyl. Ind. Jour. Agric. Sciences, 42(6): 505-508.
- Jhoopty, J.S. and Grover, R.K.(1971). Rhizoctonia root rot of cucurbit and its control in India. Indian Phytopath. 24(3): 571-574.

- Joffe, A.Z.(1969). The mycoflora of fresh and stored Ground-nut Kernalns in Israel, Mycopath. Mycol. appl. 39(3-4): 255-264.
- Kaiser, W.J. (1972a). Occurrence of three fungal diseases of chickpea in Iran. FAO Pl. Prot. Bull. 20: 73-78.
- Kaiser, W.J., Okhovar, M. and Massahebi, G.H.(1973). Effect of seed treatment fungicide on control of Ascochyta rabiei in chickpea seed intercepted with the pathogen. Pl. Dis. Reprtr. 57(9): 742-746.
- Kamal and Verma, A.K.(1979 publ. 1980). Seed-borne mycoflora of arhar (T-21), effect of culture filtrate of some isolates on seed germination and fungicidal seed treatment. Ind. Jour. Mycol. & Plant Path. 9(1): 41-45.
- Kanapathipillai, V.S.(1982). Seed mycoflora of hyacinth beans (Lablab niger) and long beans (Vigna sesquipedalis) and their pathogenic importance. Trans. Brit. Mycol. Soc. 78(43): 503-508.
- Kanipathipillai, V.S. and Hashim, Z.B.(1981). Seed-borne fungi of Brassica chinensis L. and Brassica rapa and their pathogenic importance. Malay. Agric. Jour. 53(2): 90-96.
- Kararup H.A. and Rivera, S.C.(1982). Effect of two fungicides and two insecticides on the germination and emergence of pea (Pisum sativum L.). Agro. Sur. 10(2): 75-78.
- Karmkar, S., Subudhi, S. and Badiddyopadhyay, B.(1980). Studies ,on seed mycoflora of soybean. Ind. Jour. Microbiology, 20(3): 236-238.
- Karkova,A.P., Shirko, V.N. and Nikitina, K.V.(1974). Micro-flora and its role in the decrease of seed germination of pepper and eggplant. Trudy po Prikladnoi Botanike, Genetike i Selektssii, 51(3): 224-236.

- Kaul, J.L.(1972). Efficacy of various treatments on the viability and microflora of onion (Allium cepa) seeds in storage. Ind. Jour. Mycol. & Plant Path. 2(1): 35-39.
- Karwarsa, S.S. and Mohnder Singh (1982). Seed mycoflora of cluster bean in Haryana and their control by seed treatments. Indian Phytopath. 35(3): 501-502.
- Kaul, J.L.(1973). Comparative effect of long term storage after various treatments on the viability and Mycoflora of bean (Phaseolus vulgaris) Seeds. Ind. Jour. Mycol. & Plant Path. 3(1): 17-20.
- Keenan, John, G., Moore,Howard, D., Oshma; Nagayoshi and Jenkins, Leonard E.(1974). Effect of bean root rot on dry land pinto bean production in South Western Colorado. Pl. Dis. Reprtr. 58: 890-892.
- Kerr, Eric D. and Steadman, James R.(1974). White mold damage to Western Nebraska dry beans. Annual Report of the Bean Improvement Cooperative, 17: 50-51.
- Khandelwal, G.L. and Prasad, R.(1968). Seed mycoflora of watermelon (Citrullus lanatus Thumb.) Bull. Grain Technol. 6(4): 207-223.
- Khandelwal, G.L. and Prasada,R.(1970). Seed mycoflora of cucumber (Cucumis sativus) Indian Phytopath., 23(1): 37-43.
- Khandelwal,G.L. and Prasada, R.(1970). Seed mycoflora of Bottle gourd, Labdev Jnl. Sci. Technol., 8-B(1):44-46.
- Khare, M.N.(1975). Studies on wilt of lentil. Annual Report 1974-1975. Jabalpur, M.P., India: Jawaharlal Nehru Krishi Vishwavidyalaya, 29 pp.

- Kirik, N.N. and Steblyuk, N.I.(1972). Effectiveness of Orthocide against Fusarium root rot of pea. Khinniya v. sel'skorn Khozyaistve, 10(10): 47-48.
- Linkovskaya, I.K.(1976). Effects and after effects of Zineb and heat treatments of seed on the incidence of disease and yield of cabbage. Byulletin Vsesoyuznogo Nauchno Issledovale'skogo Instituta Zashchity Rastenii No.3831-35.
- Ksendzova, E.N.(1974). Stimulation of phytoalexin production in pea by thiram. Byulletin Vsesoyuznogo Nauchno-Issledovatel'skogo Instituta Zashchity Rastenii No.29: 63-65.
- Kolev, E., Boyadzhiev, Kh. and Molnar, I.(1977). Semi wet mechanical disinfection of vegetable seeds. Gradinarska i lozrka Nauks, 14(5): 91-98.
- Kommendahl, Thor; Christensen, J.J., Culbertson, J.O. and Moore, M.B.(1955). The prevalence and importance of damaged seeds in Flax. Univ. Minnesota. Agr. Exp. Sta. Tech. Bull. 215: 400.
- Momoto, Y. and Kimura, T.(1982). Studies on the mycoflora associated with cucurbit seeds grown in Kumaun-Almora region. Madras Agric.Jour., 69(8): 548-550.
- Kotova, V.V.(1977). Biological basis and the development of control measures against disease of leguminous crop in the non-cheronozum zone. Treidy Vsesoyuznogo Nauchno-Issledovatel'skogo Instituta Zashchity Rastenii, No.53, 62-66.
- Kudrina, V.N.(1967). Change in microflora and germination of vegetable seeds during storage. Izv. timiryazev. Sel. Khoz. Akad., 2: 175-183.

- Kuhne, W.(1983). Seed dressing for better seedling emergence. Nachrichtenblatt fiir den Pflanzen der DDR 37(5): 101-102.
- Kuhn, J.(1857). Uber das Vorkommen von Anguillula an erkrankten Blaukopfen von Dipsacus fullonum L. Zeitschr. Wiss. Zool. 9: 129-37.
- Kumar, K. and Nene, Y.L.(1969). Antifungal properties of Cleome isocandra L. extracts. Indian Phytopath., 21(4): 445.
- Kumari, V. and Karan, D.(1981). Seed mycoflora of cowpea (Vigna cotjahg) and its effect on germination. Ind. Jour. Botany, 4(2): 187-190.
- Kuniyasu, K. and Nakamura, H.(1978). Seed transmission of Fusarium wilt of bottle gourd, Lagenaria siceraria used as root stock for water melons IV. The use of dry heating as a practical means of disinfection. Bulletin of the Vegetable and Oruamental crops Research Station. A.No.4, 149-162.
- Kushi, K.K. and Khare, M.N.(1979). Seed-borne fungi of sesame (Sesamum indicum) and their significance. Seed Research, 7(1): 48-53.
- Kutova, I. and Filipova, N.(1977). Effect of seed treatment with some trace elements and antibiotics of Pseudomonas lachrymans on cucumber. B"lg. Plod., Zelench i. konservi No.3, 30-32.
- Kuz'mina, G.(1966). Antibiotic against wilt. Zashch. Rast., Mosk, 11(4): 31.
- Lafon, R. and Bugaret, T.(1968). Treatment of onion seeds against white rot. Phytiat.Phytopharm, 17(2): 67-72.

- Lafon, R. and Bugaret, T.(1970). Effectiveness of benomyl against white rot of Garlic and Onion. Pytiat. Phytopharm. 19(1): 3-7.
- Lambat, A.K., Siddiqui, M.R., Nath, R.,Majumdar, A. and Rani, I.(1974). Seed-borne fungi of sugarbeet in India with special reference to Phoma betae trunk and its control. Seed Research, 2: 33-40.
- Lambhate, S.S. and Bhide, V.P.(1976). Some fungi from pea seeds and their effect on germination. Jour. Maharashtra Agric. Univ., 1(1): 35-36.
- Lasca, C.C.(1978). Studies on the mycoflora of bean (Phaseolus vulgaris L.) seeds. Biologica, 44(6):125-134.
- Leach, J.G.(1940). "Insect transmission of plant diseases". Mc. Graw-Hill Book Company, Inc. U.S.A.
- Lewis, J.A. and Lumsden, R.D.(1984). Reduction of pre-emergence damping-off of peas caused by Pythium ultimum with calcium oxide. Canad.Jour. Plant Path. 6(3): 227-232.
- Lifshitz, R., Lifshitz, S. and Baker, R.(1985). Decrease in incidence of Rhizoctonia pre-emergence damping-off by use of intergrated chemical and biological controls. Plant Disease, 69(5): 431-434.
- Lo,S.L.(1977). Control of imported vegetable seed-borne organism by antibiotics. Plant Protection Bulletin Taiwan, 19(1): 55-60.
- Lopez, Rosa, J.H.(1969). Phoma spp., the causal agent of Pigeon pea canker. Phytopathology, 59,(10): 1347-1349.

- Lorenzini, G., Mannerucci, G.F., Triolo, E. and Gambogi, P. (1975). Some pathogenic aspects of Fusarium oxysporum f. sp. pisi isolated from pea seed and efficacy of treatment with systemic fungicides. Rivista di Patologia vegetale IV, 11(112): 25-50.
- Lumsden, Robert, D., Adams, Peter B., Abawi, George and Steadman, James R.(1974). Sclerotinia workshop-Conference, 30th October - 1st November 1974, Beltsville Maryland, 3p.
- Lutynska, Regina (1968). Investigation on diseases of seed onion caused by Botrytis spp. in the vegetable growing region of the cracow district. Acta. mycol., Warszawa, 4(1): 3-22.
- Macias, W.(1970). Trials on chemical control of U.cephulae on Onion. Biul. Warzywn. Inst. Warzywn 10: 83-103.
- McClintock, J.A.(1916). Science, 44: 786.
- Madan, S., Singh, D., Mathur, S.B. and Neergaard, P.(1975). Detection and location of seed-borne inoculum of Ascochyta rabiei and its transmission in the chickpea (Cicer arietinum). Seed Science & Technology, 3(3/4): 667-681.
- Malone, I.P. and Muskett, A.E.(1964). Seed-borne fungi. Description on 77 fungi. Proc. Int. Seed. Test. Ass. 29: 1-383.
- Mani, A. and Sethi, C.L.(1984). Influence of seed treatment on seedling emergence of chickpea in presence of Meloidogyne incognita, Fusarium oxysporum f. sp. ciceri and F.solani. Ind. Jour. Nematol. 14(1):68-69.

- Masurat, G., Paschel, R., Stephan, S. (1966). Das Auftreten der wichtigsten Krankheiten und Schädlinge der landwirtschaftlichen und gartenrischen Kulturpflanzen im Jahre 1965 im Bereich der Deutschen Demokratischen Republik. Nacher Bl. dt. Pflanzenschutzd. (Berlin) 20: 121-142.
- Maude, R.B. (1966). Testing steam/air mixture for control of Ascochyta pisi and Mycosphaerella pinodes on pea seeds. Plant Path. 15(4): 187-189.
- Maude, R.B., Presly, A.H. and Dudley, C.L. (1973). 23rd Annual Report for 1972, National Vegetable Research Station, Wellesbourne, Warwick.
- Maude, R.B. and Kyle, Ann M. (1970). Seed treatments with benomyl and other fungicides for the control of Ascochyta pisi on peas. Ann. appl. Biol. 66(1): 37-41.
- Maude, R.B. and Humpherson-Jones, F.M. (1977). Fungal diseases- 93-102 of 27th Annual Report for 1976. National vegetable Research centre. Wellesbourne, Warwick, UK (1977) 157 pp.
- Maude, R.B. and Humpherson-Jones, F.M. (1980). Studies on the seed-borne phases of dark leaf spot (Alternaria brassicicola) and grey leaf spot (Alternaria brassicae) of brassicas. Ann. appl. Biol. 95(3): 311-319.
- Maude, R.B. and Presley, A.H. (1977a). Neck rot (Botrytis allii) of bulb onion I. Seed-borne infection and its relationship to the disease in the onion crop. Ann. appl. Biol. 86(2): 163-180.
- Maude, R.B. and Presley, A.H. (1977b). Neck rot (Botrytis allii) of bulb onions II. Seed-borne infection in relationship to the disease in store and the effect of seed treatment. Ann. appl. Biol. 86(2): 181-188.

- Maude, R.B., Spencer, A., Brocklehurst, P.A. Gott, K.A. and Bainbridge, J.M.(1985). The biology and control of Alternaria dauci (leaf blight) on carrot seeds. In 34th Annual Report for 1984, National Vegetable Research Station, Wellesbourne, Warwick, U.K.
- Maude, R.B., Vizer, Ann, S. and Shurin Catriona, G.(1969). The control of fungal seed-borne diseases by means of thiram seed soak. Ann. appl. Biol. 64(2): 245-257.
- Micheli, P.A.(1723). Relazione dell'erba dettada Botanic Grobanche aerolgarment Succiamelle, fiamme emald. Occhio (S.A.R.P. Tartiini, Horence, 47 p.).
- Mills, J.T. and Bollen, G.T.(1976). Microflora of heat damaged rape-seed. Canad. Jour. Botany, 54(24): 2893-2902.
- Mirkova, E.(1976). Lab. method for detecting and determining infection of carrot seeds by fungal pathogen. Rastitelna Zashchita, 24(1): 45-47.
- Mirkova, E.(1979). Carrot seet treatment against some fungal pathogens. Gradinarska i Lozarska Nauka, 16(5/6): 92-96.
- Morall, R.A.A. and Mc Kenzie D.L.(1974). A note on the inadvertent introduction to North America of Ascochyta rabiei, a destructive pathogen of chickpea. Pl. Dis. Repr. 58: 342-345.
- Muskett, A.E.(1950). Seed-borne fungi and their significance. Trans. Brit. Mycol. Soc. 33: 1-22.
- Mukerwar, P.M., Nath, R., Lambat, A.K., Kapoor, U., Khetrapal, R.K. and Rani, I.(1980). Interception Peronospora manshurica in imported seeds of soybean. Seed Research 8(2): 170-173.

- Narain, A. and Panigrahi, C.(1971 publ. 1972). Efficacy of some fungicidal compounds to control Colletotrichum capsici in vitro and in vivo. Indian Phytopath. 24(3): 593-596.
- Narayanaswami, P. and Jagannathan, T.(1975). Seed transmission of black gram leaf crinkle virus. Phytopathologische Zeitschrift, 82(2): 107-110.
- Natti, J.J.(1966). Evaluation of seed treatments with PCNB for the control of damping-off of Table Beet Seedlings. Pl. Dis. Reprtr. 50(8): 614-617.
- Nedumaran, S. and Vidhyasekharan, P.(1981). Control of Fusarium semitectum infection in tomato seed. Seed Research, 9(1): 28-31.
- Needham, T.(1745). New Microscopical discoveries (F.Needham. Halborn England, 126).
- Neergaard, Paul (1977). Seed Pathology Vol.I. The Macmillan Press Ltd.London and Basingstoke.
- Numic, R.M.,Hauzer-Alfarevic',O., Milisevic, S.(1979). The effect of an antifungal antibiotic of the monosccharide type on Pythium debaryanum Hess. Zastita Bilja, 30(3): 283-288.
- Nene, Y.L., Thapliyan, P.N. and Krishna, Kumar (1968). Screening of some plant extracts for antifungal properties. Labdev Jnl. Sci. Technol. 6 B(4): 226-228.
- Ohh, S.H., King, T.H. and Kommendahl, T.(1978). Genotype effect of captan treated pea seed to Pythium ultimum. Pl. Dis. Reprtr. 62(3): 196-199.

- Okoga, D.M. and Jaffer, A.A.(1972). Studies on the efficacy of various fungicides against bean rust (Uromyces appendiculatus) (Pers.) Lev. Miscellaneous Report. Tropical Pesticides Res. Inst. Arusha Tanzania, No.805, 13 pp.
- Oliker, M.D.(1972). The effectiveness of new chemicals in the control of disease complex of cucumber. Trudy Vsesoyuznogo Nauchno-Issledovatel'skogo Institute Zashchity Rastenii, 35: 300-306.
- Ondrej, M.(1971). The incidence of fungi of the genus Ascochyta on bean and kidney bean. Ochr. Rost. 7(3): 226-229.
- Orlova, G.I., Kanonkov, P.F.and Sizova,T.P.(1982). The determination of pathogenicity of Fusarium sp. isolated from tomato seeds. Mikologiya i Fitopatologiya 16(5): 443-446.
- Orton, C.R.(1931). Seed-borne parasites. A bibliography. Agric. Exp. Sta. West. Virginia Univ. 47p.
- Panteleev, A.A., Memedov, A.Kh, Bagirova,Z.S. and Kyazimova, N.G.(1973). Antibiotics in the control of fungus disease. Zashchita Rastenii, No. 12: 23-24.
- Pathak, V.K., Mathur, S.B. and Neergaard (1978). Detection of Peronospora manshurica (Naum) Syd. in seeds of soybean. EPPO, Bulletin, 8(1): 21-28.
- Peresyphkin, V.F.(1966). The use of UV radiation in the control of ascochytirosis of pea (Preliminary communication). Zakhyst, Roslyn., 3: 59-61.

- Perry, D.A. and Hegarty, T.W.(1971). Effect of captan seed dressing on carrot emergence. Plant Pathology 20(1): 29-32.
- Petkar, A.S., Utikar, P.G. and More, B.B.(1977). Control of collar rot of double bean caused by Macrophomina Phaseolina (Maubl.) Ashby. Mysore Jour. Agric. Sci. 11(1): 63-65.
- Petrov, B.(1972). Anthrocnose of bean. Rastitelna Zashchita, 20(5): 24-26.
- Pfleger, F.L. and Harman, G.E.(1975). Inability of storage fungi to invade pea embryos: evidence against phytoalexin involvement. Phytopathology, 65(5): 642-643.
- Pohoska, Irena (1963). Dry seed treatment of pea and Beet. Biul. Warzywn. Inst. Uprawy, Nawoz, Gelbozu, 6: 335-343.
- Porter, R.H.(1949). Recent development in seed technology. Bot. Rev. 15: 221-344.
- Pozhar, Z.A.(1971). Increasing seed treatment in the field and control of black leg. Sakh. Svekla 16(3): 32-34.
- Prem Nath (1976). "Vegetables for the tropical region", Published by ICAR, New Delhi.
- Presley, A.H. and Maude, R.B.(1980). The effect of seed treatment with benzimidazole based fungicides on infection of the foliage of over wintered salad onions by Botrytis cinerea. Ann. appl. Biol. 94(2): 185-196.

- Prevost, B.(1807). M'c moire sur la cause immediate de la carie ou charbon desbles et de plusieurs autres maladies des plantes, et zur les preservatifs dela carie (Translated from French by George Wannamaker ___ Keit under the title, Memoir on the immediate cause of Bunt or Smut of wheat, and of several other disease of plants, and on preventives of Bunt.) American Phytopathological society. Phytopathological classics No.6 Menasha, Wis 1939, 98p.
- Randhawa, H.S. and Aulakh, K.S.(1984). Efficacy of hot water treatment to control seed-borne fungi of raya (Brassica juncea coss.) Ind. Jour. Plant Path. 2(1): 73-76.
- Rauckyte, E.K.(1965). Action of some chemical prepratians against the causal agent of ascochyttosis of Broad Beans, A.fabae (1. Effectiveness of antibiotics. 2. Effectiveness of fungicides.) Liet. TSR Mokslu Akad. Darb. Ser. C., 1(36): 109-118.
- Reddy, M.V.(1980). Calizin, M. and effective fungicide for eradication of Ascochyta rabeiei in Chickpea seed. Int. Chickpea Nesletter No.3, 12:
- Remnant, R.(1637). A discourse of historie of bees unto is added the causes and cure of blasted wheat. And some remedies of blasted hope and rice and fruit. Together with the cause of sumtty wheat, all of which are very useful for this later age 47 p. London.
- Rosca, I.(1976). Distribution and intensity of infection of soybean seed by Peronospora manshurica Naum-syd. in Romania during 1973-1975. Probleme de Protectia Plantelor 4(1): 15-22.
- Rout, B.K. and Rath, G.C.(1972). Note on seed-borne diseases of chilli (Capsicum annuum). Indian Phytopath., 25(4): 597-598.

- Roy, K.W.(1982). Seedling disease caused in soybean by species of Colletotrichum and Glomerella. Phytopathology 72(8): 1093-1096.
- Russel, P.E. and Mussa, A.E.A.(1977). An evaluation of potential seed treatments to control Fusarium golani f.sp. phaseoli, the cause of foot and root rot of Phaseolus vulgaris. Jour. Agric. Sci. 89(1): 235-238.
- Russel, R.C.(1956). Longevity studies with wheat seed and certain seed-borne fungi (Abstr.).Proc.Canad. Phytopath. Soc. 24: 22p.
- Sackston, W.E.(1969). Sclerotium bataticola on seeds of cowpea (Vigna sinensis). Pl. Dis. Reprtr. 53(6): 438-439.
- Sanchez - Dominguez, R., Moreno Martinez, E. and Zanteno Zevada, M.(1971). Studies on the storage of soyabean seed of the variety Tropicana. Boleutin de la Societed Mexicana de Micologia. No.5: 47-55.
- Sarode, M.S. and Kadam, V.C.(1977). A seed-borne disease of brinjal (Solanum mlongena) Journal of Maharashtra Agric. Univ. 2(1): 80-82.
- Schmidt, J.L.(1955). Wheat storage research at Hutchinson, Kansam and James Town, North Dakota, U.S. Deptt. of Agric. Tech. Bull. 1113.
- Scott, D.J. and Wenham, H.T.(1973). Occurrence of two seed-borne pathogens Alternaria radicina and A.dauci on imported carrot seed in Newzealand. Newzealand Jour. Agric. Res., 16(2): 247-250.

- Segura, G. and Diaz-Polanco, C.(1975). Treatment of seed and seedlings of Phaseolus vulgaris with benomyl (benlate) to prevent infection by Rhizoctonia solani CIARCO, 5(1-4): 37-40.
- Sekhon, A.S. and Shivapuri, T.N.(1971). Fungi of certain stored vegetable seeds in relation to pre and post emergence effects. Mycopath. Mycol. appl. 43(2): 211-215.
- Sen, C., Srivastava, S.N. and Agnihotri, V.P.(1974 publ. 1975). Seedling disease of Sugar beet and their chemical control. Indian Phytopath., 27(4): 596-602.
- Sharma, Y.R. and Chohan, J.S.(1971). Control by thermotherapy of seed-borne vegetable Marrow mosaic virus. Pl. Prot. Bull. F.A.O. 19(4): 86-88.
- Sheppard, J.W. and Needham, S.N.(1980). Verticillium wilt of alfa-alfa in Canada. Occurrence of seed-borne inoculum, Canad. Jour. Plant Path. 2(3): 159-162.
- Shirsat, A.M. and Kale, U.V.(1979). Effect of fungicidal seed treatment on germination and seedling vigour of Chickpea (Cicer arietinum L.) Tropical grain legume Bulletin, No.16, 29-32.
- Shukla, D.N. and Bhargava, S.N.(1978). Evaluation of fungicides against seed-borne infections. Proc. Nat. Acad. Sci. India B. 48(4): 193-198.
- Shukla, P., Singh, R.R. and Mishra, A.N.(1979). Efficacy of seed dressing fungicides for the control of pea wilt. Pesticides. 13(4): 40-41.

- Sddiqui, M.R., Dharam Singh and Gaur, A.(1977). Prevalence of chilli anthracnose fungus on seeds and its effective control. Seed Research. 5(1): 67-72.
- Sindhan, G.S.(1984). Evaluation of fungicides against occurrence of angular leaf spot of French bean caused by Phaeoisariopsis griseola. Ind. Jour. Plant Path. 2(1): 39-42.
- Singh, U.P., Pathak, K.K., Khare, M.N. and Singh R.B.(1979). Effect of leaf extract of garlic on Fusarium oxysporum f. sp. ciceri, Sclerotinia sclerotiorum and on gram seeds. Mycologia, 71(3): 556-564.
- Singh, K., Srivastava, S.N., Sen, C. and Agnihotri, V.P.(1973 publ. 1974). Seed mycoflora of sugar beet (Beta vulgaris L.). and its fungicidal control. Proc. Indian Nat. Sci. Acad. B. 39(6): 695-700.
- Singh, D.V., Ray Chaudhuri, S.P. and Seth, M.L.(1970). Two new leaf spot diseases of brinjal (Solanum melongena L.) Current Science 39(23): 541-542.
- Singh, Tribhuwan and Singh, Dalbir (1982). Transmission of seed-borne inoculum of Macrophomina phaseolina from seed to plant. Proc. Indian Acad. Sci. Plant Sciences 91(4): 3457-370.
- Sinha, O.K. and Khare, M.N.(1977). Control of seed-borne Macrophomina phaseolina and Fusarium equiseti. by hot water treatment of cowpea seeds. Seed Research, 5(1): 20-22.
- Skowronski, Z.(1976). Aetiology and the possibilities of control of soft rot of curds of seed cauliflowers. Roczniki Nauk Rolniczych, E.5(2): 35-42.

- Sohi, H.S.(1976). Effect of seed dressers against seed-borne infection in certain vegetable crops. Madras Agric. Jour. 63(5/7): 426-427.
- Soteros, J.E.(1979). Pathogenicity and control of Alternaria radicina and A.dauci in carrots. Newzealand Jour. Agric. Res. 22(1): 191-196.
- Sowell, G.(1965). The effect of seed treatment on seed-borne pathogens of Gaur, Pl. Dis. Reprtr. 49(11): 895-897.
- Srivastava, R.N.(1984). Seed-borne fungi from Indian seed lots of Coriandrum sativum Linn. their significance and control. Jour. Ind. Botan. Soc. 63(2): 181-185.
- Srivastava, R.N. and Gupta, G.S.(1982). Location of extent of conidia and mycelium of various fungi in Verbena seed. Jour. Ind. Botan. Soc. 61(2/3): 151-155.
- Staron, T., Allard, C., Heitz, Suzuanne and Billet, Denise (1969): Purification and antifungal properties of a triterpenic heteroside extracted from the stem and leaves of A.arvensis var. phoenicea. Phytiat. Phytopharm. 18(3): 161-166.
- Suhag, L.S. and Suryanarayan, D.(1976 publ. 1977). Some aspects of seed health testing with respect to seed-borne fungi of pulse crops grown in Haryana. Ind. Jour. Mycol. & Plant Path. 6(1): 32-36.
- Tahvonen, R.(1979). Seed-borne fungi on cruciferous cultivated plants in Finland and their importance in seedling raising. Jour. Sci. Agric. Soc. Finland. 51(5): 227-279.
- Tai Luang Haun and Musa Bin Jamail (1975). Seed-borne pathogen in okra fruit rot MARDI Res. Bulletin, 3(2): 38-45.

- Tamiet, G. and Garibaldi, A.(1978). A root rot of spinach
Piedmont and Liguria. Aetiology and Control. Infor-
matore Fitopatologica, 28(9): 7-11.
- Tasca, G. and Hulea (1973).. Effect of chemical treatment
on fungi causing disease of carrot-during storage.
Revista de Horticulture Si Viticulture, 22(10):22-27.
- Taylor,J.D.(1973). 23rd Annual Report for 1972. National
Vegetable Research Station, Wellesbourne, Warwick.
- Tenne, F.D.,Prasartsee, C., Machado, C.C. and Sinclair, J.B.
(1974). Variation in germination and seed-borne
pathogens among soybean seed lots from three regions
in Illinois. Pl. Dis. Repr. 58(5): 411-413.
- Teranishi, J. and Namekata, T.(1972). Fusarium wilt on pea
(Pisum sativum L.) Biologico 38(3): 91-92.
- Thakur, V., Sharma, A.D. and Munjal,R.L.(1982). Seed mycoflora
of Raphanus sativus its pathology and control from
India. Ind. Jour. Mycol. & Plant Path. 11(2): 161-163.
- Tillet, M.(1755). Dissertation swele canseque corrupt et
noireit grains de bled danse 'pis' et sur los moyers.,
de prevenir car accidans Brussels, 1755, Translation
by H.B. Humphrey, Amer. Phytopath. Soc. Phytopathological
Classic 5: 191.
- Toskov, N., Boshnakov, P. and Tsirkov, I.(1975). The numbers
and specific composition of microorganisms stored
vegetable seeds. Gradinarska i hozarska Nauka, 12(5):
70-74.
- Tripathi, M.N. and Kaushik, C.D.(1984). Studies on the
survival of Alternaria brassicae the causal organism of
leaf spot of repeseed and mustard. Madræ Agric. Jour.
71(4): 237-241.

- Tylkowska, K.(1984). Health of carrot seeds in 1978-1980. Biuletyn Instytut Hodowli i Aklimatyzacji Roslin, No. 153, 205-217.
- Tylkowska, K.(1984). Occurrence of fungi on bean seeds reproduced in different regions of Poland. Biuletyn Instytutu Hodowli i. Aklimatyzacji Roslin No.153, 185-202.
- Tyner, L.E.(1951). Control of loose smut of barley by chemical and physical treatments. Sci. Agric. 31: 187-192.
- Utikar, P.G., Gadre, U.A. and More, B.B.(1978). Seed treatment to control Fusarium wilt of pea. Pesticides, 12(11): 29-30.
- Utyugova, T. and Levteeva, N.(1977). Effectiveness of carrot seed disinfection against Alternaria disease. Latvijas Lauksainiebas Akademija Raksti, No.136, 8-10.
- Vannacci, G.(1981). Record of Fusarium moniliforme var. Wr. & Reink on onion seeds and its pathogenicity. Phytopathologia Mediterranea, 20(2/3): 144-148.
- Vartanian, V.G. and Endo, R.M.(1985). Survival of Phytophthora infestans in seed extracted from infected tomato. Phytopathology, 75(3): 375-378.
- Vegh, I., Champion, R., Bourgeois, M. and Brunef, D.(1978). Presence of Phoma valerienellae Gindrat, Semecnik Bolay on Lamb's lettuce in France. Revue Horticole, No.184, 39-43.
- Verma, R.K. and Vyas, S.C.(1977). Effect of seed treatment with systemic fungicides in gram wilt control. Pesticides, 11(1): 20-21.

- Vishunavat, K. and Shukla, P.(1981). Screening of culture filtrates of seed mycoflora of lentil upon seed germinability and seedling length. Indian Phytopath. 34(4): 410-412.
- Vir, D. and Gaur, A.(1970). Efficacy of fungicides XI: seed disinfection in relation to Rhizoctonia bataticola on okra seeds. Pesticides 4(7): 25-26.
- Volkova, A.(1964). Influence of post harvest seed treatment on sowing qualities. Zernobob. Kul'tury 10: 10-11.
- V'rbánov, V.(1973). Trials of some fungicide in the seed treatment of sugar beet against black leg. Resteniev" dni Nauki, 10(3): 151-157.
- V'rbánov, V.(1974). Presowing seed treatment of sugar beet against black leg. Rastitelna Zashchita, 22(3): 33-34.
- V'rbánov, V., K.Slavov, and A.Slavche (1984). Fungicide treatment of coated beet seeds. Rasteniev" dni Nauki 21(2): 90-96.
- Walker, J.C.(1952). "Diseases of vegetable crops". (1st ed.) Mc Graw Hill-Book Company Inc. New York, Toronto, London.
- Wallen, V.R.(1965). Field evaluation and the importance of the Ascochyta complex on peas. Canad. Jour. Plant.Sci. 45: 27-33.
- Wan Zainum, Wan Nik and Tap, M.V.(1979). Rhizoctonia solani a seed-borne pathogen of French bean in Malaysia. Pertanika, 2(1): 11-15.
- Watanabe, T.(1972). Fungi associated with commercial kidney bean seed and their pathogenicity to young seedlings of kidney bean. Ann. Phytopath. Soc. Japan 38(2):111-116.

- Watanabe, T.(1972). Macrophomine phaseoli found in commercial kidney bean seed and in soil, and pathogenicity to kidney bean seedling. Ann. Phytopath. Soc. Japan, 38(2): 100-105
- Wells, J.M. and Merwarth, F.L.(1973). Fungicides dips for controlling decay of carrots in storage for processing. Pl. Dis. Reprtr. 57(8): 697-700.
- White, J.G., Crute, I.R. and Wynn, E.C.(1984). A seed treatment for the control of Pythium damping-off diseases and Peronospora parasitica on brassicas. Ann. appl. Biol. 104(2): 241-247.
- Williams, P.H.(1967). Occurrence of Phoma lingam on cabbage seed from Australia after treatment with hot water. Pl. Dis. Reprtr. 51(7): 566-569.
- Windles, C.E. and Kommendahl, T.(1978). Factors affecting Penicillium oxalicum as seed protectant-against seedling blight of pea. Phytopathology, 68(11): 1656-1661.
- Woolman, H.M. and Himphrey, Herry, B.(1924). Studies in the physiology or control of bunt or stinking smut of wheat U.S. Deptt. Agric. Bull. 1239: 29p.
- Wu, W.S. and Lu, J.H.(1984). Seed treatment with antagonists and chemicals to control Alternaria brassicicola. Seed Sci.& Technology, 12(3) : 851-862.
- Wu, W.S., Wu, C.H.H. and Wu, K.C.(1979). Alternaria brassicicola a destructive pathogen in seed production of cauliflower Plant. Prot. Bull. Taiwan, 21(3): 294-304.

ACKNOWLEDGEMENTS

It immensely gratifies me to record my heartfelt sense of gratitude for Late Dr. Dharendra Prakash, Reader, Department of Botany, A.M.U., Aligarh who had been supervising me during his life time. He is no more, but he leaves behind for me his infinite source of inspiring genius, his good deeds, his virtues which shall always adorn my life.

I want to extend my grateful thanks to Prof. S.K.Saxena, Chairman, Department of Botany, A.M.U., Aligarh, my supervisor who generously agreed to take up the task of guiding me. Despite his work load he lent me alot of time in critically reviewing the manuscript. His able guidance, plentiful worthy suggestions made the work possible. During the entire period of the pursuit of my studies I had the good fortune of enjoying his constant help and advice, without which this work would not have been what it is.

It is pleasurable to render my sincerest thank to Mr. M.Imran Khan, who took keen interest in comparing the manuscript and responding well with his ingrudging help at my slightest request.

I submit my humble thanks and regards to my brothers, sisters and parents in particular who first generated in me the interest for the thirst of knowledge.

Thanks are also due to Mazahir Husain for typing the manuscript in time.

Last, but not least, mentions are made to my friends and colleagues Messers Shamshad, Tariq, Ashraf, Zaki, Zaman, Haroor Perwaiz, Mujeeb, Musheer, Ali Alam, Tabreiz, Shaheen and Imtiaz, whose unceasing encouragement, moral support and multifarious co-operation, in some cases not related to my field of Botany, elevated me to the position that I exist on at present.

Md. Abul Fazal
Mohd. Abul Fazal

- Yoshii, K.(1975). Seed treatment of pea with benomyl to control of Mycosphaerella pinodes, Fitopatologia, 10(1): 41-44.
- Yu, S.H., Mathur, S.B. and Neergaard, P.(1982). Taxonomy and Pathogenecity of four seed-borne species of Alternaria from sesame. Trans. Brit. Mycol. Soc. 447-458.
- Zad, J.(1979). Mycoflora of soybean seed. Ind. Jour. Plant. Path. 15(1/4): 32-33.
- Zaidi, R.K. and Prakash, D.(1983). Studies of fungi associated with cowpea seeds. Jour. Ind. Botan. Soc., 62(4): 421-422.
- Zaumeyer, W.J. and Thomas, H.R.(1957). A monographic study of bean diseases and methods for their control. U.S. Dep. Agric. Tech. Bull. 868. Revised ed. 255p.
